

**Food consumption of the invasive amphipod *Dikerogammarus villosus* in field  
mesocosms and its effects on leaf decomposition and periphyton**

Susanne Worischka<sup>1,2</sup>, Luise Richter<sup>1</sup>, Anne Hänig<sup>1</sup>, Claudia Hellmann<sup>2,3</sup>, Jochen Becker<sup>2</sup>,  
Pavel Kratina<sup>4</sup> and Carola Winkelmann<sup>2</sup>

<sup>1</sup> Dresden University of Technology, Institute of Hydrobiology, Zellescher Weg 40, 01217  
Dresden, Germany

<sup>2</sup> University Koblenz-Landau, Institute of Integrates Natural Sciences, Universitätsstr. 1,  
56070 Koblenz, Germany

<sup>3</sup> biota – Institut für ökologische Forschung und Planung GmbH, Nebelring 15, 18246  
Bützow, Germany

<sup>4</sup> Queen Mary University of London, School of Biological and Chemical Sciences, Mile End  
Road, London E1 4NS'

Corresponding author: Susanne Worischka

Email: [susanne.worischka@tu-dresden.de](mailto:susanne.worischka@tu-dresden.de)

Phone: +49(351)46339628

Fax: +49(351)46337108

Co-authors' Email addresses:

Luise Richter: [luise.richter2@tu-dresden.de](mailto:luise.richter2@tu-dresden.de)

Anne Hänig: [haenig@googlemail.com](mailto:haenig@googlemail.com)

Claudia Hellmann: [hellmann@institut-biota.de](mailto:hellmann@institut-biota.de)

Jochen Becker: [jochenbecker@uni-koblenz.de](mailto:jochenbecker@uni-koblenz.de)

Pavel Kratina: [p.kratina@qmul.ac.uk](mailto:p.kratina@qmul.ac.uk)

Carola Winkelmann: [cawinkelmann@uni-koblenz.de](mailto:cawinkelmann@uni-koblenz.de)

## Abstract

Invasive species can affect native communities by replacing competitors, overexploiting prey species or altering ecosystem structure. One example is the Ponto-Caspian amphipod *Dikerogammarus villosus* which has established large populations in European rivers and is widely considered as a main cause for the decline of native benthic invertebrates. This effect has been mainly associated with direct predation, whereas the indirect effects via competition for primary resources are poorly understood and possibly underestimated. To assess the probability of those indirect effects, we performed five outdoor flow-through mesocosm experiments in three European rivers, manipulating the density of *D. villosus*. We quantified its in-situ food consumption during three 24-h gut content surveys in the mesocosms. Gut evacuation rates for correction were measured in the laboratory for different food sources and under continuous feeding. We analysed the invader's effects on primary resources by quantifying periphyton biomass and community leaf litter decomposition in the mesocosms at different *D. villosus* densities. The observed remarkably high food consumption rates (0.38-1.27 mg mg<sup>-1</sup> d<sup>-1</sup>, in dry mass/dry body mass) of *D. villosus* can be attributed mainly to its high gut evacuation rates. The leaf litter decomposition rates indicate that *D. villosus* is an efficient shredder; however, there was no effect on the periphyton biomass. Our results indicate that *D. villosus* may be a strong competitor with primary consumers in benthic food webs of invaded rivers, with not only direct but also indirect negative effects on benthic communities. High consumption rates together with an opportunistic feeding behaviour probably promote the invasion success of this amphipod.

**Keywords:** biofilm grazing, CPOM, daily ration, feeding activity, gut evacuation, river food web, shredder

## Introduction

The spread of invasive alien species is a global phenomenon associated with a decline of native biodiversity and changes in ecosystem structure and function (Hooper *et al.*, 2005, Strayer & Dudgeon, 2010). In European and North American river systems, particularly severe effects have been caused by various invasive species from the Ponto-Caspian region (Leuven *et al.*, 2009), among them several amphipods (Bollache *et al.*, 2004, Josens *et al.*, 2005, Palmer & Ricciardi, 2005, van Riel *et al.*, 2006). A prominent example for an invasive amphipod affecting Central European river communities is *Dikerogammarus villosus* SOVINSKY. This species invaded the Rhine System in 1995 (Haas *et al.*, 2002, Leuven *et al.*, 2009) and the Elbe system via the Mittellandkanal in 1998 (Bij de Vaate *et al.*, 2002) and, more recently, spread to Southern and Western Europe (e.g. Italy in 2006, Casellato *et al.*, 2006, United Kingdom in 2010, MacNeil *et al.*, 2010). In the invaded systems, it rapidly establishes high densities and often displaces both native and previously arrived alien species (Haas *et al.*, 2002, Jazdzewski *et al.*, 2004, Koop *et al.*, 2008, Hellmann *et al.*, 2016). Aside from its large body size (maximum 30 mm), other traits such as a high fecundity probably make *D. villosus* a successful invader (Devin *et al.*, 2004, Kley & Maier, 2006, Poeckl, 2009). Its success may be also enhanced by broad tolerance towards environmental factors and/or anthropogenic stressors such as salinity and temperature (Bruijs *et al.*, 2001, Grabowski *et al.*, 2007, Bacela-Spychalska *et al.*, 2013), some pesticides (Bundschuh *et al.*, 2013) and eutrophication (Brauns *et al.*, 2007). Although being generally broad, its tolerances are not always broader than those of other amphipods (Maazouzi *et al.*, 2011, Gergs *et al.*, 2013, Poznanska *et al.*, 2013). However, *D. villosus* is often able to compensate for this by different types of behaviour, such as hiding, low locomotor activity or dominance in competition for refuges (Gabel *et al.*, 2011, Becker *et al.*, 2016, Borza *et al.*, 2017). In addition, the not only omnivorous but highly flexible and opportunistic feeding behaviour of this species can support compensation (e.g. Platvoet *et al.*, 2009b, Dodd *et al.*, 2014). The potentially negative

78 predatory impacts of *D. villosus* were the focus of numerous studies because the species  
79 displays very aggressive behaviour against other invertebrates (Dick & Platvoet, 2000, Dick  
80 *et al.*, 2002, MacNeil & Platvoet, 2005, Boets *et al.*, 2010). Predation on fish eggs was also  
81 observed in laboratory experiments (Casellato *et al.*, 2007, Taylor & Dunn, 2017). Therefore,  
82 *D. villosus* is often regarded as a predator with direct negative effects on other benthic  
83 species, especially on amphipods in invaded habitats (MacNeil *et al.*, 2011). This may result  
84 in negative effects of the *D. villosus* invasion on the ecosystem functions maintained by these  
85 prey taxa, such as coarse particulate organic matter (CPOM) decomposition. Because *D.*  
86 *villosus* has displayed low shredding efficiencies in several laboratory studies, it is regarded  
87 as unable to compensate the loss of other shredders (e.g. Piscart *et al.*, 2011, Boeker & Geist,  
88 2015, Jourdan *et al.*, 2016). On the other hand, some studies suggest this species may have  
89 similar shredding capabilities to *Gammarus roeselii* or *G. pulex* (Gergs & Rothhaupt, 2008,  
90 Bundschuh *et al.*, 2013, Truhlar *et al.*, 2014). Moreover, effects of *D. villosus* on overall leaf  
91 shredding rates depend on abiotic factors, e.g. decrease with increasing flow velocity (Felten  
92 *et al.*, 2008) or conductivity (Truhlar *et al.*, 2014). At high temperatures particularly, *D.*  
93 *villosus* seems to be a more efficient shredder than native gammarids (Truhlar *et al.*, 2014,  
94 Kenna *et al.*, 2017), although the invader's predation rate also increases with temperature  
95 (Van der Velde *et al.*, 2009). With one exception (Felten *et al.*, 2008), all the above  
96 mentioned studies comparing *D. villosus* with native amphipods were conducted under  
97 laboratory conditions.

98  
99 *D. villosus* seems to be an opportunistic feeder with a very broad diet outside the laboratory.  
100 The fatty acid composition of *D. villosus* in a French reservoir suggests that decaying  
101 terrestrial plant material (including microorganisms) constituted a significant proportion of its  
102 diet (Maazouzi *et al.*, 2007). In the River Rhine and the River Elbe, *D. villosus* has a  
103 relatively low trophic position, as indicated by its stable isotope signature, and seems to

consume plant-based resources in comparable amounts to animal prey (Hellmann *et al.*, 2015). In addition, genetic diet analysis indicates that *D. villosus* does not consume other invertebrates regularly in the River Rhine (Koester *et al.*, 2016). This is in accordance with the morphology of its mouthparts which are not specialized for a predatory life style (Mayer *et al.*, 2008) but are suited for various feeding techniques. Therefore, the omnivorous *D. villosus* might act as a predator but also as a competing primary consumer in a benthic community.

The exceptionally high growth rates and high reproduction potential of *D. villosus* (Devin *et al.*, 2004) suggest that it has high consumption rates, high assimilation efficiency or substantial energy allocation into somatic growth (Gergs & Rothhaupt, 2008, Becker *et al.*, 2016). Either way, the food consumption by extremely dense *D. villosus* populations observed in the field (Haas *et al.*, 2002, Koop *et al.*, 2008, Hellmann *et al.*, 2016) can be expected to have significant effects on resources. However, to date, food consumption rates of *D. villosus* have been only estimated in small-scale laboratory settings (e.g. Gergs & Rothhaupt, 2008, Truhlar *et al.*, 2014, Boeker & Geist, 2015, Jourdan *et al.*, 2016) rather than in the field where more realistic impacts effects on the invaded community can be directly assessed.

The subtraction method (e.g. Naylor *et al.*, 1989, Gergs & Rothhaupt, 2008) is predominately used in laboratory consumption estimations, because of its simple and time-efficient applicability under standardized conditions. The method is based on the amount of the remaining food after a (most often 24-h) feeding experiment on pre-defined and pre-weighed food sources. In contrast, the *in-situ* method (Bajkov, 1935, Elliott & Persson, 1978) is based on the temporal course of the consumer's gut fullness during 24 hours and is therefore applicable also under field conditions. Consequently, gut content analysis paints a more realistic picture of the actual food consumption under natural conditions – which can differ

from laboratory trials. Moreover, the diel feeding activity patterns are easily observed in the field, providing more detailed insights into the predatory and competitive impacts on other benthic invertebrates.

Here, we used the *in-situ* method to estimate the daily food consumption of *D. villosus* in field mesocosms across different conspecific densities. Because of the opportunistic feeding behaviour of *D. villosus*, its food consumption potentially includes CPOM (e.g. leaf litter) and biofilms (e.g. periphyton). This might make the invader an efficient exploitative competitor for benthic shredders and grazers. Therefore, we evaluated the effects of different *D. villosus* biomasses on leaf litter and periphyton in the mesocosms, i.e. under natural conditions. We tested the hypothesis that *D. villosus* would have a positive effect on the community leaf decomposition rate and a negative effect on periphyton biomass. We studied the effects in three lotic ecosystems with a different invasion history and dominance of *D. villosus*: the River Rhine (invaded 1995, low native biodiversity) and the River Elbe (invaded 2001, higher native biodiversity) in Germany, and the River Bure in the Norfolk Broads, U.K. (invaded 2012, higher native biodiversity) (MacNeil *et al.*, 2013, Hellmann *et al.*, 2016).

## **Methods**

### Field mesocosm experiments

Five mesocosm experiments were conducted in total, two in the middle section of the River Rhine (km 660, near Sankt Goar, Germany, 50.16987 N, 7.66981 E), two in the upper River Elbe (km 66, near Dresden, Germany, 51.09415 N, 13.65110 E) and one in River Bure (near Wroxham, U.K., 52.714604 N, 1.405625 E). The experiments lasted 4–5 weeks. The Rhine experiments were performed in autumn 2013 (starting Oct 17) and spring 2014 (starting May 9). In River Elbe, the experiments were performed in autumn 2012 (starting Sep 9) and spring 2015 (starting May 7). The experiment in River Bure was performed in spring 2016 (starting

April 26). Before each experiment, high-grade steel mesh baskets (20 mm mesh size, Fig. 1) were filled with natural substratum (from coarse gravel to fist-sized stones, about 20–120 mm grain size). The base area of a basket was 0.1 m<sup>2</sup> and substrate depth was approximately 0.15–0.2 m. Because this depth can be colonized by *D. villosus* in similar coarse substrates of the river bed (L. Richter, personal observations), the used substrate is comparable to field habitat conditions. The baskets were exposed to the river bed, allowing for colonization by site-specific invertebrate communities for 4–6 weeks (for community composition in the baskets, see Table S2 in the appendix). After colonization, the baskets were carefully transferred to the mesocosms (Figure 1), i.e. set into flumes which were mounted on three floating pontoons and closed on both sides with 2-mm steel mesh (except the Elbe experiment in autumn 2012: 16 mm at the upstream end). There were three flumes on each pontoon and the experiments started with eight baskets per flume; baskets were sampled without replacement. The density of *D. villosus* in the flumes was manipulated at the start of the experiment in order to obtain three density treatments (Fig. 1): natural density (reached in the baskets after colonization on the river bed), high density (twice the natural density), and low density (as near zero as possible). This was achieved by the following procedure: all baskets of each flume were very carefully emptied and re-filled (to ensure equal amount of handling) but, as far as possible, all *D. villosus* individuals from the low density treatment flume were transferred to the high density flume. During the experiments, the mesh closing the flumes was cleaned 1–3 times per week. On these occasions, environmental factors were measured, including water temperature (°C), oxygen concentration (mg L<sup>-1</sup>; multiprobe HQ40d, Hach, USA), current velocity (m s<sup>-1</sup>; Mini-Air 2, Schiltknecht, Switzerland) and light intensity (except River Bure, due to technical problems) (mmol m<sup>-2</sup>; portable quantum photometer, LI-COR, USA). Mean values and mean daily ranges between all flumes are given in Table S1 in the appendix.

## Monitoring of the benthic community in the mesocosms

Benthic invertebrate density and biomass in the mesocosm flumes were estimated one day after the manipulation (initial sample) and 4 weeks later by emptying one or two baskets from each flume and collecting all invertebrates. The benthic samples were rinsed over a 500  $\mu\text{m}$  sieve and stored in 80 % ethanol. Invertebrates were identified to the lowest possible taxonomic level (Elliot & Mann, 1998, Eggers & Martens, 2001, Glöer & Meier-Brook, 2003, Eiseler, 2005, Eiseler, 2010, Waringer & Graf, 2011), enumerated and total length excluding antennae or appendices was measured to the nearest 0.1 mm using a stage micrometer under a stereo microscope. The individual biomass for each benthic specimen (mg dry mass) in the substrate baskets was calculated from mean length using length-weight relationships (Meyer, 1989, Burgherr & Meyer, 1997, Benke *et al.*, 1999, Hellmann *et al.*, 2013, Hellmann *et al.*, 2015). However, if more than 50 individuals of a single taxon occurred in a basket, only 50 randomly chosen specimens were measured and the mean individual biomass of those specimens was assigned to the remaining specimens of this taxon. The benthic biomass of each taxon ( $\text{mg basket}^{-1}$ ) was calculated as the sum of the individual biomasses.

*D. villosus* was separated into two separate size classes (adult  $\geq 8$  mm and juvenile  $< 8$  mm) to account for possible differences in feeding behaviour. Low densities of the morphologically similar *D. haemobaphes* were found in River Elbe. An accurate discrimination from *D. villosus* was possible from 2.5 – 3 mm TL for the experienced researcher, based on the shape, length and spines of the uropods. Only reliably identified *D. villosus* individuals were included in gut content analyses. For an evaluation of competition with other potential grazers and shredders in the benthic community, feeding types were assigned to all taxa according to Tachet (2002) and the [www.freshwaterecology.info](http://www.freshwaterecology.info) Database (version 7.0, Schmidt-Kloiber & Hering, 2015). Both databases use relative affinities for the single feeding types which add up to 100% for each taxon, thus facilitating the use of mixed feeding types (Chevenet *et al.*, 1994). Each taxon with an affinity  $\geq 10\%$  for the feeding type



‘grazer’ or ‘shredder’ in the literature was assigned to that feeding type, otherwise to the feeding type ‘others’. Taxa with affinities  $\geq 10\%$  for both ‘grazer’ and ‘shredder’ were classified according the feeding type with the higher affinity value. The purpose of this procedure was to mirror rather the feeding potential of the invertebrates than their realized feeding behaviour because the actual diet composition is often very variable.

#### Estimation of the food consumption of *D. villosus*

The estimation of daily food consumption of *D. villosus* was possible in three mesocosm experiments: in the River Rhine, in spring and autumn, and the River Elbe, in spring (not in all five experiments due to logistical and experimental constraints). The daily food rations were estimated during 24-h field samplings, approximately three weeks after the start of the experiment, in the baskets of the natural-density and high-density treatment flumes. In each flume, at least five individuals were collected every 4 hrs, frozen in liquid nitrogen and transported to the laboratory, where they were stored at  $-18^{\circ}\text{C}$  until further processing. The contents of pharynx and gut (hereby referred to as gut contents) were separated from the body under a dissecting microscope. Gut contents and body tissue (the latter including the empty gut and pharynx) were placed on separate pre-weighted small glass microfiber filter cuts, freeze-dried for 20-24 hrs at  $-57^{\circ}\text{C}$  and weighed to the nearest 0.001 mg. The number of collected individuals ( $n = 5$  to 36 per time point) differed according to the total *D. villosus* abundance in the baskets. If sample size was  $\leq 5$ , each individual was weighed onto a separate filter cut and its gut content on another. If a larger number of individuals per sample was available, 2-3 individuals were pooled and weighed onto one filter cut (and their pooled gut contents on another), in order to save filter material, space and time. In the calculation of the gut fullness index, the contents were related to ‘empty’ mass  $m_e$ ; here *D. villosus* dry body

mass minus dry mass of pharynx contents and gut contents. For pooled individuals, the same was done with the pooled body tissue mass and the pooled gut content mass.

The *in-situ* daily ration of *D. villosus* was estimated from the gut fullness according to Elliott and Persson (1978), as the sum of the consumption during the 4-h sampling intervals. For each interval, the samples from natural and high density flumes were pooled because *D. villosus* biomasses of the treatments did not always differ significantly due to migration effects (C. Winkelmann, unpublished data). The gut fullness indices observed at the intervals were corrected with an exponential evacuation rate, which was estimated in laboratory experiments (Heroux & Magnan, 1996). Two such experiments were conducted, for two experimental food sources (A and B) at  $14 \pm 1^\circ\text{C}$  (see also Richter *et al.*, in press): Individuals were collected in River Elbe and acclimatized in cages in an indoor flume with stones as refuges. They were fed with willow leaves (*Salix* sp., pre-conditioned for 2 weeks in aerated river water) and live or frozen chironomid larvae. The experimental food sources A (pre-conditioned willow leaves) and B (live chironomid larvae) were provided prior to the actual experiments after a 24-h (food source A) or 12-h starvation phase (food source B). During the evacuation experiments, the individuals were removed from their experimental food source, kept in groups of 5 (A) or 3 (B) and allowed to feed continuously on a well distinguishable second food source (post-A and post-B) for each experimental food. Food source post-A were paper colour-coding dots soaked for 12 h in river water and food source post-B were pre-conditioned willow leaves. Gut content samples were taken at 7 time points (0, 1, 3, 5, 9, 16 and 24 h) starting at the time of switching from food source (A to post-A and B to post-B). The experimental conditions were kept as similar as possible to those in the mesocosms and the river, by providing a near-natural habitat structure with refuges, a slight water movement due to the aeration, a season-specific light-dark cycle of 16:8 h, and keeping the animals in

groups. The paper coding dots were used because they were eaten readily and could easily be separated from the leaves during gut analysis. Their digestibility was tested in preliminary experiments over 7 days (Richter et al., in press). Although the paper dots were evacuated more slowly than willow leaves which might have resulted in a slight underestimation of the willow leaf evacuation rate, the animals were not affected negatively. The gut evacuation rate was estimated by fitting an exponential regression to the gut content data over time for each experimental food source. The mean of the negative slopes of the two regressions ( $0.195 \pm 0.039$ , mean  $\pm$  se, for chironomid larvae and  $0.245 \pm 0.048$  for willow leaves), 0.22, was used as evacuation rate (expressed in  $\text{mg mg}^{-1} \text{h}^{-1}$ ) in the calculation of  $C_d$  (Elliott & Persson, 1978).

In order to account for the temperature dependence of food consumption, the *in-situ* daily ration was corrected for the difference between actual mean water temperature during each *in-situ* consumption experiment (Table 1) and the temperature during the evacuation rate experiments ( $14^\circ\text{C}$ ) by applying Van't Hoff's equation after solving it for the *in-situ* daily ration (Vant Hoff, 1896). We used a mean  $Q_{10}$  value of 1.74 for this correction (Becker *et al.*, 2016). The cumulative daily consumption  $C_{\text{cum}}$  was calculated in  $\text{mg dry mass m}^{-2}$  basket base area ( $\text{mg m}^{-2} \text{d}^{-1}$ ), for the period between the initial and second benthic sample of each mesocosm experiment, i.e. roughly 4 weeks. Temperature correction with a  $Q_{10}$  of 1.74 was applied, using the differences between the actual mean temperature during the consumption experiments and the mean daily temperatures during the whole period.

#### Evaluation of effects on primary resources

CPOM decomposition was measured directly as leaf decomposition rate during each mesocosm experiment in all rivers (except River Elbe in autumn). We used leaf litter bags

filled with 2.5 g pre-conditioned and dried willow leaves (*Salix* sp. from local riparian vegetation), with a mesh size of 1.5 mm and an ample window of 15.0 mm mesh on the upper side to allow invertebrate shredders to access the leaves. The bags were exposed approximately one week after the start of a mesocosm experiment on the substrate surface of every basket. They were sampled weekly for 3-4 weeks by randomly collecting and carefully emptying two bags and weighing the contents after removing all animals and drying at 50°C. Each sampled bag was marked (to avoid double sampling) and re-exposed with about 2.0 g of replacement leaves until the end of the mesocosm experiment in order to avoid affecting decomposition rate by a change in resource availability. Additionally, in two experiments (River Rhine spring; River Elbe spring), 0.2 mm mesh bags excluding macroinvertebrates were exposed and sampled in parallel to get an estimation of the microbial leaf decomposition rate. The leaf decomposition rate was calculated from the decrease of leaf dry mass over time by fitting a linearized negative exponential decay model (Benfield, 2006), for each flume separately.

As an indirect measure of community grazing in the baskets, the periphyton biomass was quantified 4 weeks after the start of each experiment (expressed in mg chlorophyll-a per cm<sup>2</sup> stone surface area). Periphyton was sampled from 2-3 stones out of the uppermost substrate layer in each sampled basket by brushing off the light-exposed surface (total sampled area 165 ± 91 cm<sup>2</sup>, mean ± SD) with tap water. The samples were frozen in liquid nitrogen and stored in the dark at -80°C until analysis. The chlorophyll-a content was measured in a defined subsample volume after freeze-drying, homogenization and subsequent ethanol extraction (Wetzel & Likens, 2000) using a luminescence spectrometer (LS 50B, Perkin-Elmer, Rodgau, Germany) at 667 nm emission wavelength. The sampled surface area of the

stones was measured for all biofilm samples by carefully wrapping in aluminium foil and weighing the foil cuts afterwards (in relation to a reference cut of 10 cm<sup>2</sup> area).

### Statistical analyses

The effect of *D. villosus* biomass on leaf decomposition rate and periphyton biomass was analysed by fitting linear mixed-effects models (Pinheiro, 2000, Bates *et al.*, 2015a) using the R package lme4 (Bates *et al.*, 2015b). This allowed the combined statistical analysis of all experiments. Thus, not only (small) differences in abiotic environmental factors between the flumes within an experiment are accounted for, but also the larger seasonal and river-related differences. *D. villosus* biomass values were square root-transformed to approximate normal distribution. For CPOM decomposition rate as response variable, *D. villosus* biomass (both size classes together, mean of the initial and 4 weeks samplings) was included in the models as fixed effects and experiment and pontoon as random effects, pontoon being nested within experiment. Two sets of models were fitted: one with a common slope of the *D. villosus* effect for all experiments (a-models) and one with a random slope, i.e. the slope was potentially influenced by the experiment (b-models). The effects on periphyton biomass (chl-a as response) were modelled separately for total *D. villosus* biomass and juveniles only (< 8 mm). Here, *D. villosus* biomass (4 weeks after start) and season were included as fixed effects because of the suspected strong seasonality of periphyton growth. Similarly, we fitted two sets of models, one with fixed and one with random slope, for the *D. villosus*-season interaction effect on periphyton. We compared all the models, including the null models without fixed effects, using Akaike's Information Criterion, AIC (Johnson & Omland, 2004) to find the optimal models. The daily rations of juvenile and adult *D. villosus* in the respective mesocosm experiments were compared using permutation tests, stratified by sampling time. All statistical analyses and graphical procedures were carried out using R (version 3.3.3, R

Core Team, 2017).

## Results

In most mesocosm experiments, *D. villosus* constituted a substantial proportion of the total benthic biomass (Fig. 2, Table S2) both in the high-density and natural-density treatments. We observed comparatively low *D. villosus* biomass only in the River Rhine in autumn and in the recently invaded River Bure. In our experimental units, non-native taxa dominated the benthic communities in the rivers Rhine (87.3 – 97.7% biomass) and Elbe (74.0 – 94.5%), in contrast to River Bure (4.6 – 15.7%). Potential grazers aside from *D. villosus*, were important in the River Elbe (in autumn, mainly the invasive isopod *Jaera sarsi*) and the River Bure (native and invasive snails) but occurred in low biomasses in the River Rhine. Potential shredders other than *D. villosus* included the invasive amphipod *Echinogammarus ischnus* (syn. *Chaetogammarus ischnus*) in the River Rhine and, in low numbers, *D. haemobaphes* in the River Elbe in spring, both species showing omnivorous feeding, which includes leaf-shredding. In the River Bure, the main native shredder was the caddis larva *Halesus radiatus*.

*D. villosus* had high *in-situ* consumption rates in all mesocosms, consuming average daily rations of 38 – 127% of the body weight (Table 1). The gut fullness index (Fig. 3) of adults and juveniles was highest in the River Rhine, spring, and lowest in the River Elbe, spring, but indicated no distinct diel pattern of feeding activity in any of the experiments. At almost all sampling times, juveniles had a slightly higher mean gut fullness index than adults, with the largest differences occurring in the evening and night hours. Applying the same evacuation rate to both size classes, the daily ration of juveniles was always higher than that of adults in the respective mesocosm experiment (permutation tests, stratified by sampling time,  $n = 41 - 176$ ,  $p < 0.01$  for all experiments).

355  
356 Considerable amounts of food were probably consumed by *D. villosus* in the baskets of the  
357 mesocosms, with maximum estimates of 11.8 g m<sup>-2</sup> basket base area. The consumption by the  
358 adults constituted the major proportion of the total daily food consumption ( $C_{cum}$ ) of *D.*  
359 *villosus* in the mesocosms in the two experiments in River Rhine (Table 1). This was due to  
360 the high proportion of adult biomass in the mesocosms (87% of the total *D. villosus* biomass  
361 in autumn and 85 % in spring). In contrast, in River Elbe, the biomass of adults and juveniles  
362 was similar (57 % adults of biomass) and the proportions in consumption nearly equal for  
363 both size classes (adults: 49% of  $C_{cum}$ ).

364  
365 Leaf decomposition rate showed an overall increase with higher *D. villosus* biomass in the  
366 four analysed mesocosm experiments (Fig. 4). The best model (based on the lowest AIC with  
367  $|\Delta AIC| \geq 2.0$  to the second best one), included *D. villosus* biomass as fixed effect and  
368 pontoon and experiment as random effects (m1a, Table 2). This indicates a significant effect  
369 of *D. villosus* biomass on CPOM decomposition in the mesocosms. The model with a variable  
370 slope for the single experiments did not describe the data more accurately than that with a  
371 uniform slope, which suggests that the underlying mechanisms of the increasing CPOM  
372 decomposition rates were similar in the experiments despite different rivers and seasons. The  
373 CPOM decomposition rate in the fine mesh bags in the natural density treatment flumes was  
374 generally lower than in the coarse mesh bags of the same flumes. This indicates that  
375 macroinvertebrates accounted for a part of CPOM decomposition in the mesocosms, although  
376 their importance seemed to differ between the experiments. In River Elbe, spring, at a high  
377 density of potential shredders, the CPOM decomposition in the fine mesh bags was much  
378 lower (0.019 g d<sup>-1</sup> compared to  $0.052 \pm 0.011$  g d<sup>-1</sup> mean  $\pm$  sd). In River Bure, at a low  
379 shredder density, it was only slightly lower in the fine mesh bags ( $0.011$  g d<sup>-1</sup> compared with  
380  $0.016 \pm 0.011$  g d<sup>-1</sup>). The periphyton (chl-a) showed no clear relationship to either total or

juvenile *D. villosus* biomass (Fig. 5). None of the models was better than the respective null model; in fact, all models were very similar according to AIC (Table3). Although periphyton biomass was mostly higher in spring than in autumn, the effect of season was also not significant according to the model selection.

## Discussion

The impact of invasive species on the trophic structure and function of communities is often negative but seems to be context-dependent (Kratina *et al.*, 2014, Jackson *et al.*, 2017). This might apply also to the omnivorous *D. villosus*, as in our field mesocosm study, we observed that the invasive *D. villosus* is a remarkably strong consumer in the Central European Rivers Elbe and Rhine. Its ability to ingest more food than its own body weight per day in field mesocosms exceeded expectations from laboratory-based experiments (MacNeil *et al.*, 2011, own unpublished data, Maier *et al.*, 2011, Truhlar *et al.*, 2014). Although *D. villosus* does use periphyton and leaf litter in the field to considerable proportions (Hellmann *et al.*, 2016, Koester *et al.*, 2016), we found no effect on periphyton biomass in any of the five experiments. However, our hypothesis postulating positive effects of *D. villosus* on leaf litter decomposition was supported by the data from four mesocosm experiments in three rivers with different benthic communities. This might be explained by the fact that *D. villosus* was an important or even the dominant shredder in terms of biomass in River Elbe. Even when other invertebrate shredders are present (in the River Rhine, previously invaded species) *D. villosus* can be an efficient shredder in river ecosystems, enhancing the community leaf litter recycling. In the systems studied here, there were nearly no native gammarids and low densities of other native shredders. This precluded a test of the common assumption that the invasion of *D. villosus* negatively affects ecosystem functioning (i.e. leaf litter decomposition) due to the replacement of (more efficient) native shredders (e.g. MacNeil *et al.*, 2011, Jourdan *et al.*, 2016). Our findings indicate that there may be exceptions from this assumption,



considering the high feeding potential of *D. villosus* and depending on the community and other environmental factors. For instance, when comparing leaf shredding rates of *D. villosus* and native amphipods, the larger body size of *D. villosus* (Kenna *et al.*, 2017) and its higher feeding efficiency at higher temperatures (Truhlar *et al.*, 2014) should be taken into account. Furthermore, it seems that intraguild predation is not always as important in the field as indicated by laboratory observations (reviewed in Jackson *et al.*, 2017). The assumption of *D. villosus* feeding substantially on CPOM was supported also by our observation that many individuals were found in and on the leaf bags, particularly in the coarse mesh area on the upper side that was obviously not suitable as a refuge because it was exposed to light. Some bags had extremely high *D. villosus* densities, which might have even dampened the biomass effects on leaf decomposition rate due to spatial interference competition in the high-density treatments. *D. villosus* is able to shred leaves due to the morphology of its mouthparts (Mayer *et al.*, 2008) and CPOM is a valuable enough food source, in particular in combination with the adhering biofilm of fungi and bacteria containing essential fatty acids (Maazouzi *et al.*, 2007, Maazouzi *et al.*, 2009). Even if animal prey can be expected to be assimilated more easily,, it is conceivable that the opportunistic and flexible feeder *D. villosus* used the easily available CPOM. Selecting the most abundant or consistently available food resource, even if it is not the energetically most profitable resource (per weight unit), can be a successful foraging strategy for some consumers (Real, 1990, Worischka *et al.*, 2015). Therefore, the relative impact of *D. villosus* is likely to depend on the community structure as well as the availability of different food sources.

The daily food consumption by the total *D. villosus* population reached maximum values of more than 10 g dry mass m<sup>-2</sup> basket area in the field mesocosms due to the high feeding rate of the juveniles and the high biomass of the adults. With a dependency on any single food source and even at a mixed diet, this consumption is enormous and shows the considerable

potential of this invader as a predator or exploitative competitor. Although these values are coarse estimates, the results are quite transferrable to ‘real’ field conditions because *D. villosus* reaches densities of more than 3000 ind m<sup>-2</sup> in Central European rivers (Haas *et al.*, 2002) and can dominate macroinvertebrate communities in terms of biomass (Hellmann *et al.*, 2015). The combination of the high consumption rates and high benthic densities of this invader suggests the existence of drastic effects on resources (basal resources and/or potential prey) under natural conditions. It is therefore possible that aside from the often observed strong direct effects of *D. villosus* on other species, such as predation (e.g. Dick & Platvoet, 2000, MacNeil *et al.*, 2011) or displacement from microhabitats (e.g. Casellato *et al.*, 2008, Borza *et al.*, 2017), indirect effects by exploitation competition may also contribute to the negative consequences for invaded communities.

The large difference between our consumption estimates and the values found in other studies can be attributed mainly to methodology, i.e. the experimental conditions as well as the estimation method itself. Most estimations of the feeding rate of *D. villosus* were performed in the laboratory under highly artificial conditions, such as small experimental tanks and providing a modicum of refuge (e.g. Truhlar *et al.*, 2014, Boeker & Geist, 2015). Also the use of single individuals (Gergs & Rothhaupt, 2008, Piscart *et al.*, 2011, Jourdan *et al.*, 2016) might affect the feeding rate. A combination of more semi-natural conditions in laboratory feeding experiments, such as larger tanks with abundant refuge availability and the keeping of the animals in groups, with realistic estimations of gut evacuation rates, can result in much higher feeding rates (0.54-0.89 mg mg<sup>-1</sup> d<sup>-1</sup>, dry mass/ dry body mass, Richter *et al.*, in press) compared to the above-mentioned studies (all less than 0.4 mg mg<sup>-1</sup> d<sup>-1</sup>). Therefore, the higher feeding rates observed in our field mesocosms are plausible. Assuming that field experiments mirror the complex situation in river ecosystems better than laboratory assays, we suggest that

our findings improve the estimate of the potential impact of *D. villosus* in invaded European rivers.

The periodicity of feeding activity of *D. villosus* was weak in the mesocosms with average gut fullness being relatively constant but showing high between-individual variation. This is in accordance with behavioural observations from other studies, such as a high between- and within-individual variability in swimming activity (Bierbach *et al.*, 2016) and a strong affinity for refuges such as stones or pebbles (Platvoet *et al.*, 2009a, Kobak *et al.*, 2015). The latter behaviour enables the animals to feed even in the presence of predators, e.g. fish. An important assumption of the gut content method is the strong mathematical dependence of consumption rate on gut evacuation rate (Elliott & Persson, 1978, Worischka & Mehner, 1998) which, physiologically, may in turn depend on the ingestion rate (Eggers, 1977). This was observed for *Daphnia* sp. (Gillis *et al.*, 2005) but is likely to occur in many other invertebrates. Thus, the amplitude of gut fullness over time might be dampened by the fact that ingested food is evacuated more slowly when no fresh food is following. The dependence was accounted for in the consumption estimation by using an evacuation rate determined under continuous feeding. Although it is not possible to eliminate its influence on in-situ gut content, we assume periodicity of feeding activity to be of minor importance, because *D. villosus* has been observed to have no distinct diurnal activity rhythm in previous behavioural experiments (Richter *et al.*, in press; P. Lommatzsch, unpublished data). Continuous feeding of *D. villosus* over the whole day, especially of the more predatory adults, would have consequences for all potential prey animals, reducing the possibilities for predator avoidance to merely spatial segregation.

Because *D. villosus* is at least able to feed on periphyton (Platvoet *et al.*, 2009b), we analysed also potential grazing effects. However, the periphyton quantity was not influenced by *D.*

*villosus* biomass in the mesocosms. The first possible explanation is the presence of more efficient grazers in some experiments, such as snails (especially *Viviparus contectus* Millet, 1813, Table S2) in River Bure. They might have masked any *D. villosus* effects simply due to much higher biomasses and higher grazing rate. Another reason for the lack of *D. villosus* effects on periphyton could be the dominance of bottom-up effects on autotrophic periphyton (Keldsen, 1996, Sturt *et al.*, 2011) in the only slightly shaded flumes. This is supported by our observation of strong algal periphyton growth during the three spring experiments with a temporary dominance of filamentous algae especially in River Elbe. A third explanation is that *D. villosus*, especially the adults, probably used other resources such as CPOM. Juveniles, which can be assumed to have a higher proportion of algae in their diet (Platvoet *et al.*, 2006, and own, non-quantitative observations during the gut content analyses), most likely accounted for a minor part of the total consumption except in River Elbe in spring.

In conclusion, *D. villosus* is probably not only a predator but also a competitor for some basal resources in many benthic food webs and has the potential to positively affect the ecosystem function of leaf litter decomposition. The combination of high consumption rates, and omnivorous and opportunistic feeding behaviours probably contributes to the population persistence of this invader (Kratina *et al.*, 2012) and its strong potential to alter the structure and dynamics of native benthic communities.

## **Acknowledgements**

This study was part of a project supported by the German Research Foundation (DFG, Emmy Noether program, WI 3591/1). We would like to thank Christian Sodemann, Hendrik Sallinger, Wolfram Remmers, Madlen Gerke, Felix Grunicke, Victor Reinhold, Christoph Köbsch, Peggy Lommatzsch, Eike Dusi, Johannes Berger and Chien-Fan Liu for their help in the field and with sample processing. Without the logistic support of the German Waterways

and Shipping Administration (Wasser- und Schifffahrtsamt of Dresden and of St. Goar) and the Broads Authority (U.K.), this study would not have been possible. We thank Thomas Petzoldt very much for advice in statistical analysis, especially concerning mixed-effects models, and the anonymous reviewers for their constructive comments improving the manuscript substantially.

## References

- Bacela-Spychalska K, Grabowski M, Rewicz T, Konopacka A, Wattier R (2013) The 'killer shrimp' *Dikerogammarus villosus* (Crustacea, Amphipoda) invading Alpine lakes: overland transport by recreational boats and scuba-diving gear as potential entry vectors? *Aquatic Conservation-Marine and Freshwater Ecosystems* 23: 606-618, <http://dx.doi.org/10.1002/aqc.2329>
- Bajkov AD (1935) How to estimate the daily food consumption of fish under natural conditions. *Transactions of the American Fisheries Society* 65: 288-289
- Bates D, Machler M, Bolker BM, Walker SC (2015a) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67: 1-48
- Bates D, Maechler M, Bolker B, Walker S (2015b) lme4: Linear mixed-effects models using Eigen and S4. R-package version 1.1-10. <http://CRAN.R-project.org/package=lme4>. (accessed 07 Mar 2017)
- Becker J, Ortmann C, Wetzel MA, Koop JHE (2016) Metabolic activity and behavior of the invasive amphipod *Dikerogammarus villosus* and two common Central European gammarid species (*Gammarus fossarum*, *Gammarus roeselii*): Low metabolic rates may favor the invader. *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology* 191: 119-126, <http://dx.doi.org/10.1016/j.cbpa.2015.10.015>
- Benfield E (2006) Decomposition of leaf material. In: F. Hauer & G.A. Lamberti (eds), *Methods in Stream Ecology*. Elsevier, Amsterdam, London, San Diego, pp. 711-720

536 Benke AC, Huryn AD, Smock LA, Wallace JB (1999) Length-mass relationships for  
 537 freshwater macroinvertebrates in North America with particular reference to the  
 538 southeastern United States. *Journal of the North American Benthological Society* 18:  
 539 308-343, <http://dx.doi.org/10.2307/1468447>

540 Bierbach D, Laskowski KL, Brandt AL, Chen W, Jourdan J, Streit B, Plath M (2016) Highly  
 541 variable, unpredictable activity patterns in invasive, but not native amphipod species.  
 542 *Aquatic Ecology* 50: 261-271, <http://dx.doi.org/10.1007/s10452-016-9573-4>

543 Bij De Vaate A, Jazdzewski K, Ketelaars HaM, Gollasch S, Van Der Velde G (2002)  
 544 Geographical patterns in range extension of Ponto-Caspian macroinvertebrate species in  
 545 Europe. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 1159-1174,  
 546 <http://dx.doi.org/10.1139/f02-098>

547 Boeker C, Geist J (2015) Effects of invasive and indigenous amphipods on physico-chemical  
 548 and microbial properties in freshwater substrates. *Aquatic Ecology* 49: 467-480,  
 549 <http://dx.doi.org/10.1007/s10452-015-9539-y>

550 Boets P, Lock K, Messiaen M, Goethals PLM (2010) Combining data-driven methods and lab  
 551 studies to analyse the ecology of *Dikerogammarus villosus*. *Ecological Informatics* 5:  
 552 133-139, <http://dx.doi.org/10.1016/j.ecoinf.2009.12.005>

553 Bollache L, Devin S, Wattier R, Chovet M, Beisel JN, Moreteau JC, Rigaud T (2004) Rapid  
 554 range extension of the Ponto-Caspian amphipod *Dikerogammarus villosus* in France:  
 555 potential consequences. *Archiv für Hydrobiologie* 160: 57-66,  
 556 <http://dx.doi.org/10.1127/0003-9136/2004/0160-0057>

557 Borza P, Huber T, Leitner P, Remund N, Graf W (2017) Current velocity shapes co-existence  
 558 patterns among invasive *Dikerogammarus* species. *Freshwater Biology* 62: 317–328  
 559 <http://dx.doi.org/10.1111/fwbi.12869>

560 Brauns M, Garcia XF, Pusch MT, Walz N (2007) Eulittoral macroinvertebrate communities  
 561 of lowland lakes: discrimination among trophic states. *Freshwater Biology* 52: 1022-  
 562 1032, <http://dx.doi.org/10.1111/j.1365-2427.2007.01750.x>  
 563 Bruijs MCM, Kelleher B, Van Der Velde G, De Vaate AB (2001) Oxygen consumption,  
 564 temperature and salinity tolerance of the invasive amphipod *Dikerogammarus villosus*:  
 565 indicators of further dispersal via ballast water transport. *Archiv für Hydrobiologie* 152:  
 566 633-646  
 567 Bundschuh M, Gergs R, Schadt S, Schulz R (2013) Do differences in sensitivity between  
 568 native and invasive amphipods explain their coexistence in Lake Constance? A case  
 569 study with lambda-cyhalothrin. *Chemosphere* 92: 483-489,  
 570 <http://dx.doi.org/10.1016/j.chemosphere.2013.01.106>  
 571 Burgherr P, Meyer EI (1997) Regression analysis of linear body dimensions vs. dry mass in  
 572 stream macroinvertebrates. *Archiv für Hydrobiologie* 139: 101-112  
 573 Casellato S, La Piana G, Latella L, Ruffo S (2006) *Dikerogammarus villosus* (Sowinsky,  
 574 1894) (Crustacea, Amphipoda, Gammaridae) for the first time in Italy. *Italian Journal of*  
 575 *Zoology* 73: 97-104, <http://dx.doi.org/10.1080/11250000500502293>  
 576 Casellato S, Masiero L, La Piana G, Gigliotti F (2008) The alien amphipod crustacean  
 577 *Dikerogammarus villosus* in Lake Garda (N-Italy): the invasion continues. In: W.  
 578 Rabitsch, F. Essl & F. Klingenstein (ed^eds), *Biological invasions: from ecology to*  
 579 *conservation. Proceedings of the European Conference on Biological Invasions. Vienna.*  
 580 *Neobiota Journal*, pp. 115-122  
 581 Casellato S, Visentin A, La Piana G (2007) The predatory impact of *Dikerogammarus*  
 582 *villosus* on fish. In: F. Gherardi (eds), *Biological invaders in inland waters: Profiles,*  
 583 *distribution, and threats. Invading Nature - Springer Series in Invasion Ecology.*  
 584 Springer, Dordrecht, pp. 495-506

585 Chevenet F, Dolédec S, Chessel D (1994) A fuzzy coding approach for the analysis of long-  
 586 term ecological data. *Freshwater Biology* 31: 295-309, [http://dx.doi.org/10.1111/j.1365-](http://dx.doi.org/10.1111/j.1365-2427.1994.tb01742.x)  
 587 [2427.1994.tb01742.x](http://dx.doi.org/10.1111/j.1365-2427.1994.tb01742.x)  
 588 Devin S, Piscart C, Beisel JN, Moreteau JC (2004) Life history traits of the invader  
 589 *Dikerogammarus villosus* (Crustacea : Amphipoda) in the Moselle River, France.  
 590 *International Review of Hydrobiology* 89: 21-34,  
 591 <http://dx.doi.org/10.1002/iroh.200310667>  
 592 Dick JTA, Platvoet D (2000) Invading predatory crustacean *Dikerogammarus villosus*  
 593 eliminates both native and exotic species. *Proceedings of the Royal Society B-Biological*  
 594 *Sciences* 267: 977-983  
 595 Dick JTA, Platvoet D, Kelly DW (2002) Predatory impact of the freshwater invader  
 596 *Dikerogammarus villosus* (Crustacea: Amphipoda). *Canadian Journal of Fisheries and*  
 597 *Aquatic Sciences* 59: 1078-1084, <http://dx.doi.org/10.1139/F02-074>  
 598 Dodd JA, Dick JTA, Alexander ME, Macneil C, Dunn AM, Aldridge DC (2014) Predicting  
 599 the ecological impacts of a new freshwater invader: functional responses and prey  
 600 selectivity of the 'killer shrimp', *Dikerogammarus villosus*, compared to the native  
 601 *Gammarus pulex*. *Freshwater Biology* 59: 337-352, <http://dx.doi.org/10.1111/fwb.12268>  
 602 Efron B (1979) 1977 Rietz Lecture: Bootstrap methods - another look at the jackknife. *Annals*  
 603 *of Statistics* 7: 1-26, <http://dx.doi.org/10.1214/aos/1176344552>  
 604 Eggers DM (1977) Factors in interpreting data obtained by diel sampling of fish stomachs.  
 605 *Journal of the Fisheries Research Board of Canada* 34: 290-294  
 606 Eggers TO, Martens A (2001) Bestimmungsschlüssel der Süßwasser-Amphipoda (Crustacea)  
 607 Deutschlands, Erik Mauch Verlag, Dinkelscherben, Germany, 68 pp.  
 608 Eiseler B (2005) Bildbestimmungsschlüssel für die Eintagsfliegenlarven der deutschen  
 609 Mittelgebirge und des Tieflandes, Erik Mauch Verlag, Dinkelscherben, Germany, 112 pp.



610 Eiseler B (2010) Taxonomie für die Praxis. Bestimmungshilfen-Makrozoobenthos (1)  
 611 LANUV-Arbeitsblatt 14. p. 181. Landesamt für Natur, Umwelt und Verbraucherschutz  
 612 Nordrhein-Westfalen (LANUV NRW), Recklinghausen.  
 613 Elliot J, Mann K (1998) A key to the British freshwater leeches, Freshwater Biological  
 614 Association, Ambleside, UK, 72 pp.  
 615 Elliott JM, Persson L (1978) The estimation of daily rates of food consumption for fish.  
 616 *Journal of Animal Ecology* 47: 977-991, <http://dx.doi.org/10.2307/3682>  
 617 Felten V, Doledec S, Statzner B (2008) Coexistence of an invasive and a native gammarid  
 618 across an experimental flow gradient: flow-refuge use, mortality, and leaf-litter decay.  
 619 *Fundamental and Applied Limnology* 172: 37-48, [http://dx.doi.org/10.1127/1863-](http://dx.doi.org/10.1127/1863-9135/2008/0172-0037)  
 620 [9135/2008/0172-0037](http://dx.doi.org/10.1127/1863-9135/2008/0172-0037)  
 621 Gabel F, Pusch MT, Breyer P, Burmester V, Walz N, Garcia XF (2011) Differential effect of  
 622 wave stress on the physiology and behaviour of native versus non-native benthic  
 623 invertebrates. *Biological Invasions* 13: 1843-1853, [http://dx.doi.org/10.1007/s10530-011-](http://dx.doi.org/10.1007/s10530-011-0003-1)  
 624 [0003-1](http://dx.doi.org/10.1007/s10530-011-0003-1)  
 625 Gergs R, Rothhaupt KO (2008) Feeding rates, assimilation efficiencies and growth of two  
 626 amphipod species on biodeposited material from zebra mussels. *Freshwater Biology* 53:  
 627 2494-2503, <http://dx.doi.org/10.1111/j.1365-2427.2008.02077.x>  
 628 Gergs R, Schlag L, Rothhaupt KO (2013) Different ammonia tolerances may facilitate spatial  
 629 coexistence of *Gammarus roeselii* and the strong invader *Dikerogammarus villosus*.  
 630 *Biological Invasions* 15: 1783-1793, <http://dx.doi.org/10.1007/s10530-013-0408-0>  
 631 Gillis PL, Chow-Fraser P, Ranville JF, Ross PE, Wood CM (2005) *Daphnia* need to be gut-  
 632 cleared too: the effect of exposure to and ingestion of metal-contaminated sediment on  
 633 the gut-clearance patterns of *D. magna*. *Aquatic Toxicology* 71: 143-154,  
 634 <http://dx.doi.org/10.1016/j.aquatox.2004.10.016>

635 Glöer P, Meier-Brook C (2003) Süßwassermollusken. Ein Bestimmungsschlüssel für die  
636 Bundesrepublik Deutschland, creative printing, Hamburg, Germany, 103 pp.

637 Grabowski M, Bacela K, Konopacka A (2007) How to be an invasive gammarid (Amphipoda:  
638 Gammaroidea) - comparison of life history traits. *Hydrobiologia* 590: 75-84,  
639 <http://dx.doi.org/10.1007/s10750-007-0759-6>

640 Haas G, Brunke M, Streit B (2002) Fast Turnover in Dominance of Exotic Species in the  
641 Rhine River Determines Biodiversity and Ecosystem Function: An Affair Between  
642 Amphipods and Mussels. In: E. Leppäkoski, S. Gollasch & S. Olenin (eds), *Invasive*  
643 *Aquatic Species of Europe. Distribution, Impacts and Management*. Springer  
644 Netherlands, pp. 426-432

645 Hellmann C, Schöll F, Worischka S, Becker J, Winkelmann C (2016) River-specific effects of  
646 the invasive amphipod *Dikerogammarus villosus* (Crustacea: Amphipoda) on benthic  
647 communities. *Biological Invasions* 19: 381–398, [http://dx.doi.org/10.1007/s10530-016-](http://dx.doi.org/10.1007/s10530-016-1286-z)  
648 [1286-z](http://dx.doi.org/10.1007/s10530-016-1286-z)

649 Hellmann C, Wissel B, Winkelmann C (2013) Omnivores as seasonally important predators  
650 in a stream food web. *Freshwater Science* 32: 548-562, [http://dx.doi.org/10.1899/12-](http://dx.doi.org/10.1899/12-020.1)  
651 [020.1](http://dx.doi.org/10.1899/12-020.1)

652 Hellmann C, Worischka S, Mehler E, Becker J, Gergs R, Winkelmann C (2015) The trophic  
653 function of *Dikerogammarus villosus* (Sowinsky, 1894) in invaded rivers: a case study in  
654 the Elbe and Rhine *Aquatic Invasions* 10: 385–397,  
655 <http://dx.doi.org/10.3391/ai.2015.10.4.03>

656 Heroux D, Magnan P (1996) In situ determination of food daily ration in fish: review and  
657 field evaluation. *Environmental Biology of Fishes* 46: 61-74

658 Hooper DU, Chapin FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM,  
659 Loreau M, Naeem S, Schmid B, Setälä H, Symstad AJ, Vandermeer J, Wardle DA (2005)

660 Effects of biodiversity on ecosystem functioning: A consensus of current knowledge.  
 661 *Ecological Monographs* 75: 3-35, <http://dx.doi.org/10.1890/04-0922>  
 662 Jackson MC, Wasserman BA, Grey J, Ricciardi A, Dick JTA, Alexander ME (2017) Chapter  
 663 two: Novel and disrupted trophic links following invasion in freshwater ecosystems.  
 664 *Networks of Invasion: Empirical Evidence and Case Studies* 57: 55-97,  
 665 <http://dx.doi.org/10.1016/bs.aecr.2016.10.006>  
 666 Jazdzewski K, Konopacka A, Grabowski M (2004) Recent drastic changes in the gammarid  
 667 fauna (Crustacea, Amphipoda) of the Vistula River deltaic system in Poland caused by  
 668 alien invaders. *Diversity and Distributions* 10: 81-87, [http://dx.doi.org/10.1111/j.1366-](http://dx.doi.org/10.1111/j.1366-9516.2004.00062.x)  
 669 [9516.2004.00062.x](http://dx.doi.org/10.1111/j.1366-9516.2004.00062.x)  
 670 Johnson JB, Omland KS (2004) Model selection in ecology and evolution. *Trends in Ecology*  
 671 *and Evolution* 19: 101-108, <http://dx.doi.org/10.1016/j.tree.2003.10.013>  
 672 Josens G, De Vaate AB, Usseglio-Polatera P, Cammaerts R, Cherot F, Grisez F, Verboonen  
 673 P, Vander Bossche JP (2005) Native and exotic Amphipoda and other Peracarida in the  
 674 River Meuse: new assemblages emerge from a fast changing fauna. *Hydrobiologia* 542:  
 675 203-220, <http://dx.doi.org/10.1007/s10750-004-8930-9>  
 676 Jourdan J, Westerwald B, Kiechle A, Chen W, Streit B, Klaus S, Oetken M, Plath M (2016)  
 677 Pronounced species turnover, but no functional equivalence in leaf consumption of  
 678 invasive amphipods in the river Rhine. *Biological Invasions* 18: 763-774,  
 679 <http://dx.doi.org/10.1007/s10530-015-1046-5>  
 680 Keldsen K (1996) Regulation of algal biomass in a small lowland stream: Field experiments  
 681 on the role of invertebrate grazing, phosphorus and irradiance. *Freshwater Biology* 36:  
 682 535-546  
 683 Kenna D, Fincham WNW, Dunn AM, Brown LE, Hassall C (2017) Antagonistic effects of  
 684 biological invasion and environmental warming on detritus processing in freshwater  
 685 ecosystems. *Oecologia* 183: 875-886, <http://dx.doi.org/10.1007/s00442-016-3796-x>

686 Kley A, Maier G (2006) Reproductive characteristics of invasive gammarids in the Rhine-  
687 Maine-Danube catchment, South Germany. *Limnologica* 36: 79-90,  
688 <http://dx.doi.org/10.1016/j.limno.2006.01.002>

689 Kobak J, Jermacz L, Dzierzynska-Bialonczyk A (2015) Substratum preferences of the  
690 invasive killer shrimp *Dikerogammarus villosus*. *Journal of Zoology* 297: 66-76,  
691 <http://dx.doi.org/10.1111/jzo.12252>

692 Koester M, Bayer B, Gergs R (2016) Is *Dikerogammarus villosus* (Crustacea, Gammaridae) a  
693 'killer shrimp' in the River Rhine system? *Hydrobiologia* 768: 299-313,  
694 <http://dx.doi.org/10.1007/s10750-015-2558-9>

695 Koop JHE, Schaffer M, Ortmann C, Winkelmann C (2008) Towards environmental  
696 assessment of river ecosystems by analyzing energy reserves of aquatic invertebrates.  
697 *Limnologica* 38: 378-387, <http://dx.doi.org/10.1016/j.limno.2008.05.004>

698 Kratina P, Lecraw RM, Ingram T, Anholt BR (2012) Stability and persistence of food webs  
699 with omnivory: Is there a general pattern? *Ecosphere* 3: UNSP 50,  
700 <http://dx.doi.org/10.1890/es12-00121.1>

701 Kratina P, Mac Nally R, Kimmerer WJ, Thomson JR, Winder M (2014) Human-induced  
702 biotic invasions and changes in plankton interaction networks. *Journal of Applied*  
703 *Ecology* 51: 1066-1074, <http://dx.doi.org/10.1111/1365-2664.12266>

704 Leuven RSEW, Van Der Velde G, Baijens I, Snijders J, Van Der Zwart C, Lenders HJR, De  
705 Vaate AB (2009) The river Rhine: a global highway for dispersal of aquatic invasive  
706 species. *Biological Invasions* 11: 1989-2008, [http://dx.doi.org/10.1007/s10530-009-9491-](http://dx.doi.org/10.1007/s10530-009-9491-7)  
707 [7](http://dx.doi.org/10.1007/s10530-009-9491-7)

708 Maazouzi C, Masson G, Izquierdo MS, Pihan JC (2007) Fatty acid composition of the  
709 amphipod *Dikerogammarus villosus*: Feeding strategies and trophic links. *Comparative*  
710 *Biochemistry and Physiology a-Molecular & Integrative Physiology* 147: 868-875,  
711 <http://dx.doi.org/10.1016/j.cbpa.2007.02.010>

712 Maazouzi C, Piscart C, Legier F, Hervant F (2011) Ecophysiological responses to temperature  
 713 of the "killer shrimp" *Dikerogammarus villosus*: Is the invader really stronger than the  
 714 native *Gammarus pulex*? *Comparative Biochemistry and Physiology a-Molecular &*  
 715 *Integrative Physiology* 159: 268-274, <http://dx.doi.org/10.1016/j.cbpa.2011.03.019>

716 Maazouzi C, Piscart C, Pihan J-C, Masson G (2009) Effect of habitat-related resources on  
 717 fatty acid composition and body weight of the invasive *Dikerogammarus villosus* in an  
 718 artificial reservoir. *Fundamental and Applied Limnology* 175: 327-338,  
 719 <http://dx.doi.org/10.1127/1863-9135/2009/0175-0327>

720 Macneil C, Boets P, Lock K, Goethals PLM (2013) Potential effects of the invasive 'killer  
 721 shrimp' (*Dikerogammarus villosus*) on macroinvertebrate assemblages and biomonitoring  
 722 indices. *Freshwater Biology* 58: 171-182, <http://dx.doi.org/10.1111/fwb.12048>

723 Macneil C, Dick JTA, Platvoet D, Briffa M (2011) Direct and indirect effects of species  
 724 displacements: an invading freshwater amphipod can disrupt leaf-litter processing and  
 725 shredder efficiency. *Journal of the North American Benthological Society* 30: 38-48,  
 726 <http://dx.doi.org/10.1899/10-056.1>

727 Macneil C, Platvoet D (2005) The predatory impact of the freshwater invader  
 728 *Dikerogammarus villosus* on native *Gammarus pulex* (Crustacea : Amphipoda);  
 729 influences of differential microdistribution and food resources. *Journal of Zoology* 267:  
 730 31-38, <http://dx.doi.org/10.1017/s0952836905007351>

731 Macneil C, Platvoet D, Dick JTA, Fielding N, Constable A, Hall N, Aldridge D, Renals T,  
 732 Diamond M (2010) The Ponto-Caspian 'killer shrimp', *Dikerogammarus villosus*  
 733 (Sowinsky, 1894), invades the British Isles. *Aquatic Invasions* 5: 441-445,  
 734 <http://dx.doi.org/10.3391/ai.2010.5.4.15>

735 Maier G, Kley A, Schank Y, Maier M, Mayer G, Waloszek D (2011) Density and temperature  
 736 dependent feeding rates in an established and an alien freshwater gammarid fed on

737 chironomid larvae. *Journal of Limnology* 70: 123-128, <http://dx.doi.org/10.3274/jl11-70->  
738 [1-14](http://dx.doi.org/10.3274/jl11-70-1-14)

739 Mayer G, Maier G, Maas A, Waloszek D (2008) Mouthparts of the ponto-caspian invader  
740 *Dikerogammarus villosus* (Amphipoda : Pontogammaridae). *Journal of Crustacean*  
741 *Biology* 28: 1-15, <http://dx.doi.org/10.1651/07-2867r.1>

742 Meyer EI (1989) The relationship between body length parameters and dry mass in running  
743 water invertebrates. *Archiv für Hydrobiologie* 117: 191-103

744 Naylor C, Maltby L, Calow P (1989) Scope for growth in *Gammarus pulex*, a fresh-water  
745 benthic detritivore. *Hydrobiologia* 188: 517-523, <http://dx.doi.org/10.1007/bf00027819>

746 Palmer ME, Ricciardi A (2005) Community interactions affecting the relative abundances of  
747 native and invasive amphipods in the St. Lawrence River. *Canadian Journal of Fisheries*  
748 *and Aquatic Sciences* 62: 1111-1118, <http://dx.doi.org/10.1139/F05-012>

749 Pinheiro J, Bates, Douglas (2000) Mixed-Effects Models in S and S-PLUS, Springer, New  
750 York, Berlin, Heidelberg, 537 pp.

751 Piscart C, Mermillod-Blondin F, Maazouzi C, Merigoux S, Marmonier P (2011) Potential  
752 impact of invasive amphipods on leaf litter recycling in aquatic ecosystems. *Biological*  
753 *Invasions* 13: 2861-2868, <http://dx.doi.org/10.1007/s10530-011-9969-y>

754 Platvoet D, Dick JTA, Konijnendijk N, Van Der Velde G (2006) Feeding on micro-algae in  
755 the invasive Ponto-Caspian amphipod *Dikerogammarus villosus* (Sowinsky, 1894).  
756 *Aquatic Ecology* 40: 237-245, <http://dx.doi.org/10.1007/s10452-005-9028-9>

757 Platvoet D, Dick JTA, Macneil C, Van Riel MC, Van Der Velde G (2009a) Invader-invader  
758 interactions in relation to environmental heterogeneity leads to zonation of two invasive  
759 amphipods, *Dikerogammarus villosus* (Sowinsky) and *Gammarus tigrinus* Sexton:  
760 amphipod pilot species project (AMPIS) report 6. *Biological Invasions* 11: 2085-2093,  
761 <http://dx.doi.org/10.1007/s10530-009-9488-2>

762 Platvoet D, Van Der Velde G, Dick JTA, Li SQ (2009b) Flexible omnivory in  
 763 *Dikerogammarus villosus* (Sowinsky, 1894) (Amphipoda) - Amphipod Pilot Species  
 764 Project (Ampis) Report 5. *Crustaceana* 82: 703-720,  
 765 <http://dx.doi.org/10.1163/156854009x423201>  
 766 Poeckl M (2009) Success of the invasive Ponto-Caspian amphipod *Dikerogammarus villosus*  
 767 by life history traits and reproductive capacity. *Biological Invasions* 11: 2021-2041,  
 768 <http://dx.doi.org/10.1007/s10530-009-9485-5>  
 769 Poznanska M, Kakareko T, Krzyzynski M, Kobak J (2013) Effect of substratum drying on the  
 770 survival and migrations of Ponto-Caspian and native gammarids (Crustacea:  
 771 Amphipoda). *Hydrobiologia* 700: 47-59, <http://dx.doi.org/10.1007/s10750-012-1218-6>  
 772 R Core Team (2017) R: A language and environment for statistical computing. version 3.3.3.  
 773 <http://www.R-project.org/>. (accessed 07 Mar 2017)  
 774 Real LA (1990) Predator switching and the interpretation of animal choice behavior: the case  
 775 for constrained optimization. In: R.N. Hughes (eds), *Behavioural Mechanisms of Food*  
 776 *Selection*. Series G: Ecological Sciences. Springer Verlag, Berlin Heidelberg, pp. 1-22  
 777 Richter L, Schwenkmezger L, Becker J, Winkelmann C, Hellmann C, Worischka S (in press)  
 778 The very hungry amphipod: the invasive *Dikerogammarus villosus* shows high  
 779 consumption rates for two food sources and independent of predator cues. *Biological*  
 780 *Invasions*, <http://dx.doi.org/10.1007/s10530-017-1629-4>  
 781 Schmidt-Kloiber A, Hering D (2015) [www.freshwaterecology.info](http://www.freshwaterecology.info) - An online tool that  
 782 unifies, standardises and codifies more than 20,000 European freshwater organisms and  
 783 their ecological preferences. *Ecological Indicators* 53: 271-282,  
 784 <http://dx.doi.org/10.1016/j.ecolind.2015.02.007>  
 785 Strayer DL, Dudgeon D (2010) Freshwater biodiversity conservation: recent progress and  
 786 future challenges. *Journal of the North American Benthological Society* 29: 344-358,  
 787 <http://dx.doi.org/10.1899/08-171.1>

788 Sturt MM, Jansen MaK, Harrison SSC (2011) Invertebrate grazing and riparian shade as  
 789 controllers of nuisance algae in a eutrophic river. *Freshwater Biology* 56: 2580-2593,  
 790 <http://dx.doi.org/10.1111/j.1365-2427.2011.02684.x>

791 Tachet H, Richoux P, Bournaud M, Usseglio-Polatera P (2002) Invertebrés d'eau douce -  
 792 systématique, biologie, écologie. p. 587. CNRS Editions, Paris.

793 Taylor NG, Dunn AM (2017) Size matters: predation of fish eggs and larvae by native and  
 794 invasive amphipods. *Biological Invasions* 19: 89-107, [http://dx.doi.org/10.1007/s10530-](http://dx.doi.org/10.1007/s10530-016-1265-4)  
 795 [016-1265-4](http://dx.doi.org/10.1007/s10530-016-1265-4)

796 Truhlar AM, Dodd JA, Aldridge DC (2014) Differential leaf-litter processing by native  
 797 (*Gammarus pulex*) and invasive (*Dikerogammarus villosus*) freshwater crustaceans under  
 798 environmental extremes. *Aquatic Conservation-Marine and Freshwater Ecosystems* 24:  
 799 56-65, <http://dx.doi.org/10.1002/Aqc.2375>

800 Van Der Velde G, Leuven RSEW, Platvoet D, Bacela K, Huijbregts MaJ, Hendriks HWM,  
 801 Kruijt D (2009) Environmental and morphological factors influencing predatory  
 802 behaviour by invasive non-indigenous gammaridean species. *Biological Invasions* 11:  
 803 2043-2054, <http://dx.doi.org/DOI> 10.1007/s10530-009-9500-x

804 Van Riel MC, Van Der Velde G, De Vaate AB (2006) To conquer and persist: colonization  
 805 and population development of the Ponto-Caspian amphipods *Dikerogammarus villosus*  
 806 and *Chelicorophium curvispinum* on bare stone substrate in the main channel of the River  
 807 Rhine. *Archiv für Hydrobiologie* 166: 23-39, [http://dx.doi.org/10.1127/003-](http://dx.doi.org/10.1127/003-9136/2006/0166-0023)  
 808 [9136/2006/0166-0023](http://dx.doi.org/10.1127/003-9136/2006/0166-0023)

809 Vant Hoff JH (1896) Studies in Chemical Dynamics, F. Muller, Amsterdam, pp.

810 Waringer J, Graf W (2011) Atlas of Central European Trichoptera Larvae, Erik Mauch  
 811 Verlag, Dinkelscherben, Germany, 468 pp.

812 Wetzel RG, Likens GE (2000) Limnological analyses, Springer, New York, pp.



813 Worischka S, Mehner T (1998) Comparison of field-based and indirect estimates of daily  
814 food consumption in larval perch and zander. *J. Fish Biology* 53: 1050-1059  
815 Worischka S, Schmidt SI, Hellmann C, Winkelmann C (2015) Selective predation by  
816 benthivorous fish on stream macroinvertebrates – the role of prey traits and prey  
817 abundance. *Limnologica - Ecology and Management of Inland Waters* 52: 41-50,  
818 <http://dx.doi.org/10.1016/j.limno.2015.03.004>  
819  
820

## Tables


**Table 1** Daily food consumption of *D. villosus* in the mesocosm experiments.  $C_d$  and  $C_{d,Q10}$  = daily ration in dry mass per dry body mass without and with temperature correction, TL = mean total length of the analysed individuals, n = total number of samples (all samples of a 24-h survey, number in brackets = total number of individuals if they were pooled for part of the samples),  $T_w$  = water temperature during 24-h survey,  $C_{cum}$  = cumulative daily consumption based on mean *D. villosus* biomass, Prop.  $C_{cum}$  = proportion of size class in total cumulative daily consumption, aut = autumn, spr = spring. The standard error of  $C_d$  was calculated using a bootstrap procedure (Efron, 1979).

Experiment	size class	$C_d \pm se$ (g g <sup>-1</sup> d <sup>-1</sup> )	$C_{d,Q10}$	TL $\pm$ sd (mm)	n	$T_w \pm sd$ (°C)	$C_{cum} \pm sd$ (g m <sup>-2</sup> d <sup>-1</sup> )	Prop. $C_{cum}$ (%)
Rhine, aut	adult	0.43±0.02	0.38	10.92±1.61	110	11.8±0.1	1.0±0.7	81.5
	juvenile	0.64±0.06	0.56	6.12±1.00	54			18.5
Rhine, spr	adult	0.71±0.03	0.95	11.48±1.52	42 (126)	19.1±0.8	11.8±5.0	84.0
	juvenile	0.96±0.07	1.27	4.87±0.86	85			16.0
Elbe, spr	adult	0.36±0.03	0.46	9.15±1.20	122 (168)	18.6±1.1	8.5±5.1	55.5
	juvenile	0.51±0.03	0.65	5.39±1.76	185 (265)			44.5


**Table 2** Model selection of linear mixed models with CPOM decomposition rate and periphyton chl-a, respectively, as dependent variables. For periphyton chl-a, one set of models was built for total *D. villosus* biomass and one for only juvenile *D. villosus* biomass as fixed effect. Dvill = *D. villosus* biomass (all size classes), Dvill<sub>juv</sub> = juvenile *D. villosus* biomass (< 8 mm), pont = pontoon, exper = experiment (pontoon always nested within experiment), Df = degrees of freedom of the model, AIC = Akaike's information criterion, LogLik = logarithm of maximum likelihood.

Model		Parameters		Df	AIC	LogLik
fixed effects		random effects				
response: CPOM decay rate						
m1b	Dvill	pont, exper (random slope)		9	-72.8*	45.4
m0b	-	pont, exper (random slope)		8	-69.0	42.5
m1a	Dvill	pont, exper		5	-80.3*	45.1
m0a	-	pont, exper		4	-74.4	41.2
response: Periphyton (chl-a)						
m3b	Dvill, season (interaction)	pont, exper (random slope)		11	198.4	-88.2
m3a	Dvill, season (interaction)	pont, exper		7	197.4	-91.7
m2a	Dvill, season	pont, exper		6	198.3	-93.1
m1ad	Dvill	pont, exper		5	196.8	-93.4
m1as	season	pont, exper		5	199.3	-94.6
m0a	-	pont, exper		4	198.5	-95.3
m3b	Dvill <sub>juv</sub> , season (interaction)	pont, exper (random slope)		11	210.2	-94.1
m3a	Dvill <sub>juv</sub> , season (interaction)	pont, exper		7	202.2	-94.1
m2a	Dvill <sub>juv</sub> , season	pont, exper		6	201.2	-94.6
m1ad	Dvill <sub>juv</sub>	pont, exper		5	200.5	-95.2
m1as	season	pont, exper		5	199.3	-94.6
m0a	-	pont, exper		4	198.5	-95.3


\* Likelihood ratio test:  $p < 0.05$ ,  $df = 1$ ,  $\chi^2 = 4.64$  resp. 4.01




844  
845 **Figure 1** (A) Mesocosm in the river, (B) substrate basket before exposure on the river bed for  
846 colonization, (C) schematic drawing (top view) of a mesocosm with three flumes containing  
847 eight colonized substrate baskets each. The baskets were open at the top during the  
848 experiments. The three density treatments were achieved by manipulating the *D. villosus*  
849 density. Grey arrows indicate the flow of water through the flumes which were closed with  
850 2-mm steel mesh at the prow and stern ends. (D) Schematic drawing (cross section) of a  
851 mesocosm. (E) Position of the mesocosms in the river (100-200 m apart from each other) and  
852 distribution of the treatments in each mesocosm during experiments. Drawings are not to  
853 scale.



**Figure 2** Biomass of benthic invertebrates, grouped by feeding type, in the mesocosm flumes with the three *D. villosus* density treatments (mean of 0- and 4-week sampling except Elbe, autumn: only 4-week sampling). Dvill = *D. villosus* (not included in any of the three feeding types but regarded separately), ad  $\geq$  8 mm, juv < 8 mm, gra = grazer, shr = shredder, oth = others. inv = invasive or non-native taxa. aut = autumn, spr = spring. For detailed community composition see Table S2 in the appendix.




**Figure 3** Gut fullness index of *D. villosus* in the mesocosms, measured in mg gut contents mg<sup>-1</sup> empty body mass (adult, > 8 mm, black squares, and juveniles, < 8 mm, grey circles): (A) River Rhine, autumn 2013 at  $11.8 \pm 0.1^\circ\text{C}$  water temperature,  $n = 5 - 27$  per time point, (B) River Rhine, spring 2014 at  $19.1 \pm 0.8^\circ\text{C}$ ,  $n = 6 - 12$  per time point, (C) River Elbe, spring 2015 at  $18.6 \pm 1.1^\circ\text{C}$ ,  $n = 7 - 35$  per time point. All values in dry mass per dry body mass. Time corresponds to CET in (a) and to CEST in (b) and (c), grey areas mark the dark periods between sunset and sunrise.



874

875 **Figure 4** Leaf decomposition rate (mg day<sup>-1</sup>) and *D. villosus* biomass (mg basket<sup>-1</sup>; all size  
876 classes, dry mass, mean of start and 4-week sample of each mesocosm experiment). Colours  
877 indicate the mesocosm experiments and symbols (squares, triangles and circles) indicate the  
878 three pontoons used in each experiment, with three mesocosm flumes each. Regression lines:  
879 linear mixed-effects model with residuals (dashed). The regression lines are curved to account  
880 for the square-root transformation of biomass. For model specifications see text.

881



**Figure 5** Autotrophic biofilm (chl-a) and *D. villosus* biomass (all size classes, dry mass) in the mesocosms, sampled 4 weeks after start. Colours indicate the mesocosm experiments and symbols (squares, triangles and circles) indicate the three pontoons used in each experiment, with three flumes each. Linear mixed-effects model showed no fixed effects of *D. villosus* biomass or season. For model specifications see text.